

Morphometric considerations of articular surfaces of calcaneum, occurrence of enthesophytes and evolutionary significance of cuboidal facet

Prakash KG^{1,*}, Viveka S²

¹Professor & Head, ²Assistant professor, Department of Anatomy, Azeezia Institute of Medical Sciences, Kollam

*Corresponding Author:

E-mail: drprakashkg@gmail.com

Abstract:

Introduction: Pattern of talar articular facets on the superior aspect of calcaneum significantly influence in causation of subtalar arthritis. Enthesophytes from calcaneum is a major cause of heel pain. Evolutionary modification of cuboidal facet on calcaneum is an important aspect of human bipedal gait. The objective of this study was to determine the incidence of calcaneal facet pattern and incidence of enthesophytes in calcaneum.

Methodology: Thirty-four dry human calcaneum were studied in Department of anatomy, Azeezia Institute of Medical Sciences, Kollam. All bones were examined for pattern of talar articular facets and classified accordingly. Meticulous examination was carried out to evaluate the incidence of enthesophytes. Morphometry of articular facet for cuboid and talar facets were carried out.

Results: Pattern I morphologic variant was the most frequently encountered type (25 out of 34). Pattern V not observed. Enthesophytes were observed in 21 (61.7%) bones, with medial predominant medial enthesophytes. One bone showed two enthesophytes. Mean cuboidal facet surface area was 386 (± 46) square mm.

Conclusion: It was noted that pattern I was the most common morphological variant. Medial tubercular enthesophytes were most common type of abnormal bony outgrowths of calcaneum. The knowledge about variations in morphology is important for all surgeries in subtalar region.

Key words: Calcaneum, Talar articular facets, Cuboidal facet, Enthesophytes

Access this article online	
Quick Response Code:	Website: www.innovativepublication.com
	DOI: 10.5958/2394-2126.2016.00016.5

Introduction

Entheses is connective tissue between tendon or ligament and bone. Enthesophytes are abnormal bony projections at the attachment of a tendon or ligament. Plantar calcaneal enthesophytes are the bony outgrowths from calcaneum.¹ Osseous outgrowths at the plantar aspect of calcaneum was first reported in 1900 and were called Kalkaneussporn (calcaneal spur).² Such outgrowths are seen in 11-16% of general population. Nearly three fourth of patients with heel-pain have calcaneal spur.^{1,3-5} Radiographs reveal a calcaneal spur in about 50% of patients, but the exact significance of this finding is uncertain.⁶ The exact cause of formation of enthesophyte is poorly understood. According to longitudinal traction hypothesis, it is the constant wear and tear resulting from pull of flexor digitorum brevis and ossification of plantar fascia.^{7,8} Calcaneal spurs are of two types: Dorsal/posterior spurs and plantar/inferior spurs. In this study, we have evaluated the incidence of enthesophytes in dry calcaneal bone.

Anterior, middle and posterior facets on the superior aspect of talus forms the joint with talus. Review of

literature suggests that posterior facet is most constant among these facets, anterior and middle shows considerable variations.^{5,9-13} Joint pathologies in subtalar region is more common in certain morphological varieties of facets.¹⁴ Such variations influence subtalar joint stability and mobility.^{4,15} During surgeries in subtalar region, operating surgeon needs to be aware of these morphological variations of calcaneal facets. This study addresses the variations of articular facets of calcaneum. Reported variations in south Indian population are minimal. The objective of this study was to evaluate the morphometry of articular facets, to find out facet variations, report the incidence of enthesophytes and to study the evolutionary significance of cuboidal facet of calcaneum.


Methodology

The study was conducted in department of Anatomy, Azeezia Institute of Medical Sciences, Kollam using 34 dry calcaneum. Bones with developmental abnormalities and pathological changes including healed fractures were excluded. Each bone was observed for pattern of articular surfaces. Talar articular surface of each bone was photographed and were categorised into five patterns. Fusion of anterior and middle talar facets as pattern I; anterior and middle separate is pattern II; absence of anterior facet is pattern III; fusion of anterior, middle and posterior is pattern IV; and fusion of middle and posterior facet is pattern V. Minimum distance between anterior and middle facets in pattern II calcanei was measured and tabulated. Surface area of talar and

cuboid facets was measured using area calculation application. Left and right side values for all the parameters considered were tabulated separately. All calcanei bones were meticulously observed with hand-held magnifying glass for tubercles and incidence of enthesophytes were recorded.

Results

Out of 34 dry calcanei studied, 18 were right and 16 were left sided. Pattern I with fused middle and posterior facet was found in 25 bones, pattern II in 7. There was one bone with absent anterior facet and one with all facets fused. There was no pattern V bone. Wherever the distance between the facets were less than 0.1 mm and were connected by a bony bridge, it was considered to be fused. Sidewise pattern distribution is shown in graph 1.



Graph 1: Incidence of morphological patterns of calcaneal talar facets, blue bars - right calcaneum, red bars - left calcaneum. n=34

The average distance between middle and posterior facet in type I calcaneum was 0.48 (± 0.02) mm. The average distance between anterior and middle facet in type II calcaneum was 0.26 (± 0.04) mm and middle and posterior facet was 0.39 (0.01) mm. The mean surface area of anterior, middle and posterior facets in type II pattern calcaneum is given in table 1. Table 2 gives mean surface area of cuboidal facet of calcaneum.

Table 1: Surface area of individual facets of calcaneum in type II pattern (n= 7, values in square mm)

Facet	Right	Left	Mean
anterior	33.0	19.7	26.3
middle	53.3	27.3	40.3
posterior	449.7	299.3	374.5

Table 2: Tabulation of Mean surface area and perimeter of cuboidal facet for calcaneum, n=34, standard deviation in parenthesis

Cuboidal facet for calcaneum	Right	Left	Mean
Area (in square mm)	451.2 (± 32)	320.3 (± 48)	386 (± 46)
Perimeter (in mm)	76.7 (± 12)	67 (± 8)	72.1 (± 9)

Medial tubercle is the largest tubercle seen on all calcanei. Though all four anterior, medial, lateral and posterior tubercles were seen on all bones, one bone had no anterior tubercle. Enthesophytes are observed in 21 (61.7%) bones. It was more on right side. All values are tabulated in table 3. Medial enthesophytes were predominant (Figure 1). One bone even had two medial enthesophytes. Spurs varied in length and morphology and the details were not considered in present study. Out of these fourteen specimens with enthesophytes, ten were type I pattern and four were pattern II.

Table 3: Incidence of enthesophytes found in the study, n=34, percentages in parenthesis

	Right	Left	Total
Medial	8 (23.5%)	4 (11.7%)	12 (35.3%)
lateral	2 (5.9%)	0	2 (5.9%)
Posterior	4 (11.7%)	3 (8.8%)	7 (20.5%)

**Figure 1: Enthesophytes documented in the study. (blue arrows)**

Discussion

In parlance to our objectives, we have determined the incidence of patterns of talar articular facets and occurrence enthesophytes. In this study, the most frequently found pattern is type I and in this pattern calcaneum, enthesophytes were encountered maximum. Our study goes according to many of the previous workers.^{4,5,9-11,15,16} Incidence of enthesophytes is considerably higher in comparison to many of the previous reports. Our results are similar to previous studies by Bassiouni¹⁷ and Banadda¹⁶ et al, where they have studied calcaneum of older people. Thus, the higher incidence of enthesophytes in this study, may be due to considering calcaneum of older people. Similar higher incidence is reported in many of MR radiographic studies.^{6,18} These enthesophytes arise from attachment of one of plantar aponeurosis and plantar muscles, like flexor digitorum brevis, abductor hallucis and adductor digiti minimi. Since many of the spurs were from medial tubercle, it is suggested that out of these structures, attachment of plantar aponeurosis and flexor digitorum brevis plays a crucial role in formation of enthesophytes.

There are reports that subtalar arthritis is most common in type I pattern.¹⁷ So, this study shows light on possible role of pattern of calcaneal articular facet in occurrence of arthritis in a particular locality. Despite many uncertainty regarding pathogenesis of enthesophytes, many of modalities of treatment has been recommended ranging from simple excision to more recently developed techniques such as endoscopic

excision.² Most of the techniques involve release of plantar aponeurosis.

Evolutionary significance: Arboreal heritage is well marked in human foot. It is shown by its primary grasping nature. It is indicated by well developed plantar aponeurosis, strong plantar ligaments and longitudinal arches. Comparison of chimpanzee and human allows us to understand the footsteps of evolution. Primates have significantly elongated calcaneus in comparison to many non-primates.¹⁹ Calcaneal elongation can be understood as a mechanism to replace loss of foot leverage that resulted with acquired grasping hallucal metatarsal joint. This has shifted the fulcrum of distal limb segment from the metatarsal heads to tarso-metatarsal joint.²⁰ The cuboid articular facet is the most distinctive feature of human foot²¹. In humans, the cuboid articular facet on calcaneum is asymmetrical. Superior part of this articular facet projects more anteriorly than the inferior part (figure 2). The facet only assumes a more longitudinal orientation in relation to the ground when the anterior aspect of articular facet is raised as it would be normally arched foot²². There is a deeper concavity on the medial aspect of the facet, which articulates with the beak like projection of the cuboid. This arrangement makes calcaneum a 'keystone' in lateral longitudinal arch of foot. The complex nature of this articular facet combined with reciprocal architecture of cuboid allows distinctive lateral swing of human calcaneocuboid joint. This brings the joint into closed packed position during stance phase of the bipedal locomotor gait²³.




Figure 2: Side view of right calcaneum showing projection of superior part more anteriorly than the inferior part, black line shows the direction of projection.

In contrast, many of the other primates including chimpanzees, gorillas lack this anterolateral projection of cuboid articular facet on calcaneum²⁴. This does not permit locking of calcaneocuboid joint. Absence of such close packing during locomotion results in 'midtarsal break' in stance phase of ape locomotor cycle²³.

Conclusion

Thirty four human dry calcaneum morphometry, pattern of articular facet and enthesophytes were studied. It was noted that pattern I was the most common morphological variant. Medial tubercular enthesophytes were most common type of abnormal bony outgrowths of calcaneum. The knowledge about variations in morphology is important for all surgeries in subtalar region.

Conflict of Interest: None

Source of Support: Nil

References:

1. Thomas JL, Christensen JC, Kravitz SR, Mendicino RW, Schubert JM, Vanore JV, et al. The Diagnosis and Treatment of Heel Pain: A Clinical Practice Guideline—Revision 2010. *J Foot Ankle Surg.* 2010 May;49(3):S1–S19.
2. Tanz SS. Heel pain. *Clin Orthop.* 1963;28:169–78.
3. Koca T, Aydın A, Sezen D, Başaran H, Karaca S. Painful plantar heel spur treatment with Co-60 teletherapy: factors influencing treatment outcome. Springerplus [Internet]. 2014 Jan 10 [cited 2015 Nov 29];3. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3967731/>
4. Menz HB, Zammit GV, Landorf KB, Munteanu SE. Plantar calcaneal spurs in older people: longitudinal traction or vertical compression? *J Foot Ankle Res.* 2008 Aug 11;1(1):7.
5. Kullar JS, Randhawa GK, Kullar KK. A study of calcaneal enthesophytes (spurs) in Indian population. *Int J Appl Basic Med Res.* 2014 Sep;4(Suppl 1):S13–S16.
6. Abreu MR, Chung CB, Mendes L, Mohana-Borges A, Trudell D, Resnick D. Plantar calcaneal enthesophytes: new observations regarding sites of origin based on radiographic, MR imaging, anatomic, and paleopathologic analysis. *Skeletal Radiol.* 2003 Jan;32(1):13–21.
7. Tahririan MA, Motififard M, Tahmasebi MN, Siavashi B. Plantar fasciitis. *J Res Med Sci Off J Isfahan Univ Med Sci.* 2012 Aug;17(8):799–804.
8. Digiovanni BF, Nawoczenski DA, Malay DP, Graci PA, Williams TT, Wilding GE, et al. Plantar fascia-specific stretching exercise improves outcomes in patients with chronic plantar fasciitis. *J Bone Jt Surg.* 2006;88(8):1775–81.
9. Muthukumaravel N, Ravichandran D, Melani Rajendran S. Human calcaneal facets for the talus: patterns and clinical implications. *J Clin Diagn Res.* 2011;5(4):791–4.
10. Mini MP, Silotry N, Haritha KN. Morphological study on patterns of talar articular facets of human calcanei. *Int J Med Clin Res.* 2012;3(3):136–9.
11. Kang HS, Whang TS, Cho BP, Yang YC. Morphological study on talar articular facets of korean adult calcanei. *Korean J Anat.* 1990;23(2):247–52.
12. Madhavi C, Madhuri V, George VM, Antonisamy B. South Indian calcaneal talar facet configurations and osteoarthritic changes. *Clin Anat.* 2008;21(6):581–6.
13. Garg R, Dagal N, Kumar S, Shekhawat S. Study of patterns of talar articular facets of human calcanei and their clinical implications in population of Rajasthan. *IJBAMR.* 2013;2:643–50.
14. Niki H, Hirano T, Akiyama Y, Beppu M. Accessory Talar Facet Impingement in Pathologic Conditions of the Peritalar Region in Adults. *Foot Ankle Int.* 2014 Oct 1;35(10):1006–14.
15. Sharada R, Sneha K, Gupta C, Pai SR, Rairam GB. Non-metrical study of the pattern of talar articular facets in south Indian dry calcanei. *Surg Radiol Anat.* 2012 Feb 12;34(6):487–91.
16. Banadda BM, Gona O, Vaz R, Ndlovu DM. Calcaneal Spurs in a Black African Population. *Foot Ankle Int.* 1992 Jul 1;13(6):352–4.
17. Bassiouni M. Incidence of calcaneal spurs in osteoarthritis and rheumatoid arthritis, and in control patients. *Ann Rheum Dis.* 1965;24(5):490–3.
18. Ebraheim NA, Mekhail AO, Yeasting RA. Components of the posterior calcaneal facet: anatomic and radiologic evaluation. *Foot Ankle Int.* 1996;17(12):751–7.
19. Moyà-Solà S, Köhler M, Alba DM, Roig I. Calcaneal proportions in primates and locomotor inferences in *Anchomomys* and other Palaeogene Euprimates. *Swiss J Palaeontol.* 2011 Dec 8;131(1):147–59.
20. Hall-Craggs ECB. An osteometric study of the hind limb of the Galagidae. *J Anat.* 1965 Jan;99(Pt 1):119–26.
21. Prang TC. Calcaneal robusticity in Plio-Pleistocene hominins: Implications for locomotor diversity and phylogeny. *J Hum Evol.* 2015 Mar;80:135–46.
22. Susman RL. Evolution of the Human Foot: Evidence from Plio-Pleistocene Hominids. *Foot Ankle Int.* 1983 May 1;3(6):365–76.
23. Aiello L, Christopher Dean. *An Introduction to Human Evolutionary Anatomy.* Academic Press; 1990. 609.
24. Griffin NL, Miller CE, Schmitt D, D'Àoût K. Understanding the evolution of the windlass mechanism of the human foot from comparative anatomy: Insights, obstacles, and future directions. *Am J Phys Anthropol.* 2015 Jan 1;156(1):1–10.